arXiv:1407.0176v1 [math.AG] 1 Jul 2014

On the Abhyankar-Moh inequality

Roland D. Barrolleta, Evelia R. García Barroso, and Arkadiusz Płoski

July 2, 2014

Abstract

Abhyankar and Moh in their fundamental paper on the embeddings of the line in the plane proved an important inequality which can be stated in terms of the semigroup associated with the branch at infinity of a plane algebraic curve. In this note we study the semigroups of integers satisfying the Abhyankar-Moh inequality and give a simple proof of the Abhyankar-Moh embedding theorem.

This short note is conceived as an appendix to [GB-P]. Our aim is to study the semigroup of integers appearing in connection with the Abhyankar-Moh inequality which is the main tool in proving the famous embedding line theorem ([A-M], Main Theorem). Since the Abhyankar-Moh inequality can be stated in terms of the semigroup associated with the branch at infinity of a plane algebraic curve ([GB-P], Theorem 6.4) it is natural to consider the semigroups for which such an inequality holds. In what follows we use freely the properties of semigroups explained in [GB-P], Preliminaries, pp. 3-6.

Let $n > 0$ be an integer. A semigroup $G \subseteq \mathbf{N}$ will be called an *Abhyankar-Moh semigroup of degree* n if the n-minimal sequence of generators $\overline{b}_0 = n, \overline{b}_1, \ldots, \overline{b}_h$ of G satisfies the following conditions

(G1) Set $e_k = \gcd(\overline{b}_0, \ldots, \overline{b}_k)$ for $0 \leq k \leq h$ and $n_k = \frac{e_{k-1}}{e_k}$ for $1 \leq k \leq h$. Then $e_h = 1$ and $n_k > 1$ for $1 \leq k \leq h$.

(G2) $n_{k-1}\overline{b}_{k-1} < \overline{b}_k$ for $2 \leq k \leq h$.

2000 *Mathematics Subject Classification:* Primary 20M14; Secondary 32S05.
Key words and phrases: semigroups, sequences of divisors, Abhyankar-Moh inequality.
The second-named author was partially supported by the Spanish Projects PNMTM 2007-64007 and MTM2012-36917-C03-01.

(G3) $n_{h-1}\overline{b_h} < n^2$.

Our aim is to prove the following

Theorem 1 *Let G be an Abhyankar-Moh semigroup of degree $n > 1$ and let c be the conductor of G, that is the smallest element of G such that all integers bigger than or equal to it are in G. Then $c \leq (n-1)(n-2)$. If $\overline{b}_0 = n, \overline{b}_1, \ldots, \overline{b}_h$ is the n-minimal sequence of generators of G then $c = (n-1)(n-2)$ if and only if $\overline{b}_k = \frac{n^2}{e_{k-1}} - e_k$ for $k \in \{1, \ldots, h\}$.*

Proof. Let $\delta_0 = n$ and $\delta_k = \dfrac{n^2}{e_{k-1}} - \overline{b}_k$ for $k \in \{1, \ldots, h\}$. We have $\delta_k = \dfrac{n^2 - e_{k-1}\overline{b}_k}{e_{k-1}} \geq \dfrac{n^2 - e_{h-1}\overline{b}_h}{e_{k-1}} > 0$ and $\gcd(\delta_0, \ldots, \delta_k) = e_k$ for $k \in \{1, \ldots, h\}$. Let $\gamma = \sum_{k=1}^{h}(n_k - 1)\delta_k - \delta_0 + 1$. Since $\delta_k \geq e_k$ we get $\gamma \geq \sum_{k=1}^{h}(n_k - 1)e_k - e_0 + 1 = 0$ with equality if and only if $\delta_k = e_k$ for $k \in \{0, \ldots, h\}$.

On the other hand

$$\begin{aligned}\gamma &= \sum_{k=1}^{h}(n_k - 1)\left(\frac{n^2}{e_{k-1}} - \overline{b}_k\right) - n + 1 = \sum_{k=1}^{h}(n_k - 1)\frac{n^2}{e_{k-1}} - n + 1 - \sum_{k=1}^{h}(n_k - 1)\overline{b}_k \\ &= (n-1)^2 - \sum_{k=1}^{h}(n_k - 1)\overline{b}_k = (n-1)(n-2) - c\end{aligned}$$

since $c = \sum_{k=1}^{h}(n_k - 1)\overline{b}_k - \overline{b}_0 + 1$ by [GB-P], Proposition 1.5, *4*.

Therefore $c \leq (n-1)(n-2)$ since $\gamma \geq 0$ and $c = (n-1)(n-2)$ if and only if $\gamma = 0$ which is equivalent to $\delta_k = e_k$ that is $\dfrac{n^2}{e_{k-1}} - \overline{b}_k = e_k$. ■

Let $n > 1$ be an integer. A sequence of integers $e_0, \ldots, e_h$ will be called a *sequence of divisors of* n if e_k divides e_{k-1} for $k \in \{1, \ldots, h\}$ and $n = e_0 > e_1 > \cdots > e_{h-1} > e_h = 1$.

Now, we can give a simple description of the Abhyankar-Moh semigroups of degree $n > 1$ with $c = (n-1)(n-2)$. For any sequence of divisors $e_0, e_1, \ldots, e_h$ of n, we put

$$G(e_0, \ldots, e_h) = \mathbf{N}n + \mathbf{N}(n - e_1) + \cdots + \mathbf{N}\left(\frac{n^2}{e_{k-1}} - e_k\right) + \cdots + \mathbf{N}\left(\frac{n^2}{e_{h-1}} - 1\right).$$

Using Theorem 1 we check easily

Property 2 *A semigroup G is an Abhyankar-Moh semigroup of degree $n > 1$ with $c = (n-1)(n-2)$ if and only if $G = G(e_0, \ldots, e_h)$ where $n = e_0, e_1, \ldots, e_h$ is a sequence of divisors of n.*

Here is another application of Theorem 1

Property 3 *Let G be an Abhyankar-Moh semigroup of degree $n > 1$ with $c = (n-1)(n-2)$ and let $n' = \min(G \backslash \{0\})$. Then $\gcd(n, n') = n - n'$. In particular, $n - n'$ divides n.*

Proof. Let $\overline{b}_0 = n, \overline{b}_1, \ldots, \overline{b}_h$ be the n-minimal sequence of generators of G. The sequence $e_{k-1}\overline{b}_k$, $k \in \{1, \ldots, h\}$ being increasing we get that $e_0\overline{b}_1 \leq e_{h-1}\overline{b}_h < n^2$, that is $\overline{b}_1 < n$. Consequently $\overline{b}_1$ is the smallest term of the sequence $\overline{b}_0, \overline{b}_1, \ldots, \overline{b}_h$ and $n' = \min(G \backslash \{0\}) = \overline{b}_1$. By the second part of Theorem 1 $\overline{b}_1 = e_1(n_1 - 1) = n - e_1$ and the property follows. ■

Property 4 *Let G be an Abhyankar-Moh semigroup of degree n with $c = (n-1)(n-2)$ and let $\overline{\beta_0}, \overline{\beta_1}, \ldots, \overline{\beta_g}$ be the minimal system of generators of the semigroup G. Then $n = \overline{\beta_1}$ or $n = 2\overline{\beta_0}$.*
If $n = \overline{\beta_1}$ then $G = \mathbf{N}n + \mathbf{N}(n - \epsilon_1) + \mathbf{N}\left(\frac{n^2}{\epsilon_1} - \epsilon_1\right) + \cdots + \mathbf{N}\left(\frac{n^2}{\epsilon_{g-1}} - 1\right)$.
If $n = 2\overline{\beta_0}$ then $G = \mathbf{N}(n - \epsilon_0) + \mathbf{N}\left(\frac{n^2}{\epsilon_0} - \epsilon_1\right) + \cdots + \mathbf{N}\left(\frac{n^2}{\epsilon_{g-1}} - 1\right)$,

where $\epsilon_k = \gcd\left(\overline{\beta_0}, \overline{\beta_1}, \ldots, \overline{\beta_k}\right)$ for $k \in \{0, \ldots, g\}$.

Proof. The conditions (G1) and (G2) imply that $n \leq \overline{\beta_1}$ and $n \equiv 0 \pmod{\overline{\beta_0}}$ if $n < \beta_1$. We claim that if $n \neq \beta_1$ then $n = 2\overline{\beta_0}$. Indeed, if $n \neq \beta_1$ then $n = a\overline{\beta_0}$ for an integer $a > 0$ and $n = b(n - \overline{\beta_0})$ for an integer $b > 0$ by Property 3. Thus we get $a = (a-1)b$ which implies $a = 2$. The second part of the property follows from [GB-P], Proposition 1.5, *3.* ■

Remark 5 *Suppose that the sequence $\overline{b_0} = n, \ldots, \overline{b_1}, \ldots, \overline{b_h}$ satisfies the conditions* (G1), (G2) *and* (G3). *Then the sequence of positive integers $\delta_0 = \overline{b}_0 = n$, $\delta_k = \dfrac{n^2}{e_{k-1}} - \overline{b}_k$ for $k \in \{1, \ldots, h\}$, has the following properties*

(i) The sequence of divisors $\gcd(\delta_0, \ldots, \delta_k) = e_k$, $k \in \{1, \ldots, h\}$, is strictly decreasing, $e_h = 1$.

(ii) $\delta_1 < \delta_0$ and $\delta_k < n_{k-1}\delta_{k-1}$ for $k \in \{2, \ldots, h\}$, where $n_k = \frac{e_{k-1}}{e_k}$.

In contrast to the property $n_k\overline{b}_k \in \mathbf{N}n + \cdots + \mathbf{N}\overline{b}_{k-1}$ *of the sequences* $\overline{b}_0, \ldots, \overline{b}_h$ *satisfying the conditions* (G1) *and* (G2)*, in general, the condition* $n_k\delta_k \in \mathbf{N}n + \cdots + \mathbf{N}\delta_{k-1}$ *fails. For example* $(\overline{b}_0, \overline{b}_1, \overline{b}_2) = (6, 2, 17)$ *is an Abhyankar-Moh sequence with the corresponding sequence* $(\delta_0, \delta_1, \delta_2) = (6, 4, 1)$ *but* $n_2\delta_2 = 2 \notin \mathbf{N}\delta_0 + \mathbf{N}\delta_1 = \mathbf{N}6 + \mathbf{N}4$.

Abhyankar-Moh Embedding Line Theorem ([GB-P], Theorem 6.6). Assume that C is a rational projective irreducible curve of degree $n > 1$ with one branch at infinity and such that the center of the branch at infinity O is the unique singular point of C. Suppose that C is permissible and let n' be the multiplicity of C at O. Then $n - n'$ divides n.

Proof. It follows from [GB-P], Theorem 6.4 that Γ_O is an Abhyankar-Moh semigroup of degree n. Let c be the conductor of the semigroup Γ_O. Using the Noether formula for the genus of projective plane curve we get $c = (n-1)(n-2)$. Then the theorem follows from Property 3. ■

References

[A-M] Abhyankar, S.S.; Moh, T.T. Embeddings of the line in the plane. *J. reine angew. Math.* **276** (1975), 148-166.

[Ba] Barrolleta, R. Semigrupos en teoría de singularidades: aplicaciones a los códigos. *Memoria fin de Máster, Universidad de La Laguna, Tenerife* (Julio 2013).

[GB-P] García Barroso, E. R., Płoski, A. An approach to plane algebroid branches. arXiv:1208.0913. 4 Aug 2012.

Roland David Barrolleta
Departament d'Enginyeria de la Informació i de les Comunicacions
Universitat Autònoma de Barcelona
08193 - Bellaterra - Cerdanyola, Barcelona, Spain
e-mail: rolanddavid.barrolleta@deic.uab.cat

Evelia Rosa García Barroso
Departamento de Matemáticas, Estadística e I.O.
Sección de Matemáticas, Universidad de La Laguna
38271 La Laguna, Tenerife, España
e-mail: ergarcia@ull.es

Arkadiusz Płoski
Department of Mathematics
Kielce University of Technology
Al. 1000 L PP7
25-314 Kielce, Poland
e-mail: matap@tu.kielce.pl